

<b>Table S1. Different scoring systems for Kawasaki disease*.</b>		
<b>Egami Risk Scores</b>		
A cut-off point $\geq 3$ identifies patients with Kawasaki syndrome at high risk of being resistant to intravenous immunoglobulin.		
Age	<6 months	1 point
Days of illness	<4	1 point
Platelet count	< $300 \times 10^9/L$	1 point
C-reactive protein	>8 mg/dL	1 point
Alanine aminotransferase	>80 IU/L	2 points
<b>Kobayashi Risk Scores</b>		
A cut-off point $\geq 4$ identifies patients with Kawasaki syndrome at high risk of being resistant to intravenous immunoglobulin.		
Sodium	$\leq 133$ mmol/L	2 points
Days of illness at initial treatment	$\leq 4$	2 points
Aspartate aminotransferase	$\geq 100$ IU/L	2 points
Percentage of neutrophils	$\geq 80\%$	2 points
C-reactive protein	$\geq 10$ mg/dL	1 point
Age	$\leq 12$ months	1 point
Platelet count	$\leq 300 \times 10^9/L$	1 point
<b>Sano Risk Scores</b>		
Non-responsiveness to intravenous immunoglobulin is predicted by the presence of at least 2 of 3 predictors in the child with Kawasaki syndrome.		
C-reactive protein	$\geq 7.0$ mg/dL	
Total bilirubin	$\geq 0.9$ mg/dL	
Aspartate aminotransferase	$\geq 200$ IU/L	
<b>Harada scores</b>		
The Presence of at Least 4 of these 7 Criteria prompted the Administration of Intravenous Immunoglobulin to the Child with Kawasaki Syndrome.		
White blood cell count	$> 12 \times 10^3/mm^3$	
Platelet count	$< 35 \times 10^4/mm^3$	
C-reactive protein	$> 3+$	
Hematocrit	$< 35\%$	
Albumin	$< 3.5$ g/dL	
Age	$< 12$ months	
Sex	Male	
<b>Nakano scores</b>		
Low-risk group is composed of patients whose total score is 0 or positive, while high-risk group is composed of patients whose total score is negative.		
Age of onset Under 1 yr 1-2 yr Over 2 yr	Weight of Category (-0.5) Weight of Category (+0.0) Weight of Category (+0.3)	Score (-1) Score (0) Score (+1)
C-reactive protein 0 to 1+ 2+ to 4+ 5+ 6+	Weight of Category (+0.9) Weight of Category +0.4 Weight of Category (-0,0) Weight of Category (-1.9)	Score (+2) Score (+1) Score (0) Score (-3)
Platelet count Under $30 \times 10^4$ Over $30 \times 10^4$	Weight of Category (-0.4) Weight of Category (+0.2)	Score (-1) Score (+1)
*Rigante D, Andreozzi L, Fastiggi M, Bracci B, Natale MF, Esposito S. Critical Overview of the Risk Scoring Systems to Predict Non-Responsiveness to Intravenous Immunoglobulin in Kawasaki Syndrome.		